

3 Punktionstechniken und Verschlusssysteme

- [1] Schnyder G et al. Common femoral artery anatomy is influenced by demographics and comorbidity: implications for cardiac and peripheral invasive studies. *Cathet Cardiovasc Intervent* 2001; 53:289–295
- [1a] Stone PA, Campbell JE, AbuRahma AF. Femoral pseudoaneurysms after percutaneous access. *J Vasc Surg* 2014; 60(5): 1359–1366
- [2] Dotter CT, Rösch J, Robinson M. Fluoroscopic guidance in femoral artery puncture. *Radiology* 1978; 127:266–7
- [3] Bashore TM et al. Cardiac catheterization laboratory standards: a report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents (ACC/SCA & I Committee to Develop an Expert Consensus Document on Cardiac Catheterization Laboratory Standards). *J Am Coll Cardiol* 2001; 37:2170–2214
- [4] Kahlert P et al. A modified „preclosure technique“ after percutaneous aortic valve replacement. *Catheter Cardiovasc Interv* 2008; 72:877–84
- [5] Petroglou D, Didagelos M, Chalikias G et al. Manual versus mechanical compression of the radial artery after transradial coronary angiography: the MEMORY Multicenter Randomized Trial. *JACC Cardiovasc Interv* 2018; 11:1050–1058
- [6] Kwok CS, Kontopantelis E, Kinnaird T et al. British Cardiovascular Intervention Society (BCIS) and National Institute of Cardiovascular Outcomes Research (NICOR). Retroperitoneal hemorrhage after percutaneous coronary intervention: incidence, determinants, and outcomes as recorded by the British Cardiovascular Intervention Society. *Circ Cardiovasc Interv* 2018; 11(2):e005866
- [7] Eggebrecht H et al. Impact of gender on femoral access complications secondary to application of a collagen-based vascular closure device. *J Invasive Cardiol* 2004; 16:247–250
- [8] Amer O, Binger S, Desch S et al. Incidence, predictors, and treatment options of critical limb ischaemia after use of collagen plug-based vascular closure devices. *Euro-Intervention* 2015; 11:816–823
- [9] Leibundgut G, Pache J, Schulz S et al. Collagen plug vascular closure devices and reduced risk of bleeding with bivalirudin versus heparin plus abciximab in patients undergoing percutaneous coronary intervention for non st-segment elevation myocardial infarction. *J Interv Cardiol* 2013; 26:623–629
- [10] Rao AK, Pratt C, Berke A et al. Thrombolysis in myocardial infarction (TIMI) trial – phase I: hemorrhagic manifestations and changes in plasma fibrinogen and the fibrinolytic system in patients treated with recombinant tissue plasminogen activator and streptokinase. *J Am Coll Cardiol* 1988; 11:1–11
- [11] The GUSTO investigators. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. *N Engl J Med* 1993; 329:673–682
- [12] Serebruany VL, Atar D. Assessment of bleeding events in clinical trials – proposal of a new classification. *Am HJ Cardiol* 2007; 99:288–290
- [13] Mehran R, Rao SV, Bhatt DL et al. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation* 2011; 123:2736–2747
- [14] Chan YC, Morales JP, Reidy JF et al. Management of spontaneous and iatrogenic retroperitoneal haemorrhage: conservative management, endovascular intervention or open surgery? *Int J Clin Pract* 2008; 62:1604–1613
- [15] Chatterjee T, Do DD, Kaufmann U et al. Ultrasound-guided compression repair for treatment of femoral artery pseudoaneu-

- rysm: acute and follow-up results. *Cathet Cardiovasc Diagn* 1996; 38:335–340
- [16] Webber GW, Jang J, Gustavson S et al. Contemporary management of postcatheterization pseudoaneurysms. *Circulation* 2007; 115:266–274
 - [17] Omoigui NA, Califf RM, Pieper K et al. Peripheral vascular complications in the Coronary Angioplasty Versus Excisional Atherectomy Trial (CAVEAT-I). *J Am Coll Cardiol* 1995; 26:922–930
 - [18] Tisi PV, Callam MJ. Surgery versus non-surgical treatment for femoral pseudoaneurysm. *Cochrane Database Syst Rev* 2006; 25(1):CD004981
 - [19] Tisi PV, Callam MJ. Treatment for femoral pseudoaneurysms. *Cochrane Database Syst Rev* 2009; 5(2)
 - [20] Schaub F, Theiss W, Busch R et al. Management of 219 consecutive cases of postcatheterization pseudoaneurysm. *J Am Coll Cardiol* 1997; 30(3):670–675
 - [21] Schaub F, Theiss W, Heinz M et al. New aspects in ultrasound-guided compression repair of postcatheterization femoral artery injuries. *Circulation* 1994; 90(4):1861–1865
 - [22] Waksman R, King SB III, Douglas JS et al. Predictors of groin complications after balloon and new-device coronary intervention. *Am J Cardiol* 1995; 75(14):886–889
 - [23] Lumsden AB, Miller JM, Kosinski AS et al. A prospective evaluation of surgically treated groin complications following percutaneous cardiac procedures. *Am Surg* 1994; 60(2):132–137
 - [24] Zahn R, Thoma S, Fromm E et al. Do 5-F catheters reduce the incidence of a pseudoaneurysm? *Int Angiol* 1996; 15(3):257–260
 - [25] Kent KC, McArdle CR, Kennedy B et al. Accuracy of clinical examination in the evaluation of femoral false aneurysm and arteriovenous fistula. *Cardiovasc Surg* 1993; 1(5):504–507
 - [26] Hanson JM, Atri M, Power N. Ultrasound guided thrombin injection of iatrogenic groin pseudoaneurysm: Doppler features and technical tips. *Radiology* 2008; 127(1):154–163
 - [27] Kang SS, Labropoulos N, Mansour MA et al. Percutaneous ultrasound guided thrombin injection: a new method for treating postcatheterization femoral pseudoaneurysms. *J Vasc Surg* 1998; 27(6):1032–1038
 - [28] Powell A, Benenati JF, Becker GJ et al. Percutaneous ultrasound-guided thrombin injection for the treatment of pseudoaneurysms. *J Am Coll Surg* 2002; 194(1 Suppl):S53–S57
 - [29] Babu SC, Piccorelli GO, Shah PM et al. Incidence and results of arterial complications among 16,350 patients undergoing cardiac catheterization. *J Vasc Surg* 1989; 10(2):113–116
 - [30] Dangas G, Mehran R, Kokolis S et al. Vascular complications after percutaneous coronary interventions following hemostasis with manual compression versus arteriotomy closure devices. *J Am Coll Cardiol* 2001; 38(3):638–641
 - [31] Roberts SR, Main D, Pinkerton J. Surgical therapy of femoral artery pseudoaneurysm after angiography. *Am J Surg* 1987; 154(6):676–680
 - [32] Jain SP, Roubin GS, Iyer SS et al. Closure of an iatrogenic femoral artery pseudoaneurysm by transcatheter coil embolization. *Cathet Cardiovasc Diagn* 1996; 39(3):317–319
 - [33] Owen RJ, Haslam PJ, Elliott ST et al. Percutaneous ablation of peripheral pseudoaneurysms using thrombin: a simple and effective solution. *Cardiovasc Intervent Radiol* 2000; 23(6):441–446
 - [33] Samal AK, White CJ, Collins TJ et al. Treatment of femoral artery pseudoaneurysm with percutaneous thrombin injection. *Catheter Cardiovasc Interv* 2001; 53(2):259–263
 - [34] Brummer U, Salcuni M, Salvati F et al. Repair of femoral postcatheterization pseudoaneurysm and arteriovenous fistula with percutaneous implantation of endovascular stent. *Nephrol Dial Transplant* 2001; 16(8):1728–1729
 - [35] Rath J, Marx R, Ganschow US et al. Formation, therapy and prevention of false aneurysm of the femoral artery following diagnostic and interventional heart catheterization. *Dtsch Med Wochenschr* 1997; 122(24):771–777
 - [36] Heis HA, Bani-Hani KE, Elheis MA et al. Postcatheterization femoral artery pseudoaneurysms: therapeutic options: A case-controlled study. *Int Surg* 2008; 87(3):214–219

- [37] Heis HA, Bani-Hani KE, Elheis MA et al. Postcatherization femoral artery pseudoaneurysms: therapeutic options: A case-controlled study. *Int Surg* 2008; 6(3):214–219
- [38] Samuels D, Orron DE, Kessler A et al. Femoral artery pseudoaneurysm: Doppler sonographic features predictive for spontaneous thrombosis. *J Clin Ultrasound* 1997; 25(9):497–500
- [39] Feld R, Patton GM, Carabasi RA et al. Treatment of iatrogenic femoral artery injuries with ultrasound-guided compression. *J Vasc Surg* 1992; 16(6):832–840
- [40] Fellmeth BD, Baron SB, Brown PR et al. Repair of postcatheterization femoral pseudoaneurysms by colour flow ultrasound guided compression. *Am Heart J* 1992; 123(2):547–551
- [41] Zahn R, Thoma S, Fromm E et al. Pseudoaneurysm after cardiac catheterization: therapeutic interventions and their sequelae: experience in 86 patients. *Cathet Cardiovasc Diagn* 1997; 0(1):9–15
- [42] Görges G, Kunz T, Kirstein M. Non-surgical therapy of iatrogenic false aneurysms. *Dtsch Med Wochenschr* 2003; 128(1–2):36–40
- [43] Brophy DP, Sheiman RG, Amatulle P et al. Iatrogenic femoral pseudoaneurysms: thrombin injection after failed US-guided compression. *Radiology* 2000; 214(1):278–282
- [44] Fellmeth BD, Roberts AC, Bookstein JJ et al. Postangiographic femoral artery injuries: nonsurgical repair with US-guided compression. *Radiology* 1991; 178(3):671–675
- [45] Schaub F, Theiss W, Heinz M et al. New aspects in ultrasound-guided compression repair of postcatheterization femoral artery injuries. *Circulation* 1994; 90(4):1861–1865
- [46] Vazquez V, Reus M, Pinero A et al. Human thrombin for treatment of pseudoaneurysms: comparison of bovine and human thrombin sonogram-guided injection. *AJR Am J Roentgenol* 2005; 184:1665–1671
- [47] Rath J, Marx R, Ganschow US et al. Formation, therapy and prevention of false aneurysm of the femoral artery following diagnostic and interventional heart catheterization. *Dtsch Med Wochenschr* 1997; 122(24):771–777
- [48] Carroll JF, Moskowitz KA, Edwards NM et al. Immunologic assessment of patients treated with bovine fibrin as a hemostatic agent. *Thromb Haemost* 1996; 76(6):925–931
- [49] Görges G, Kunz T. Thrombin injection for treatment of false aneurysms after failed compression therapy in patients on full-dose antiplatelet and heparin therapy. *Catheter Cardiovasc Interv* 2003; 58(4):505–509
- [50] Görges G, Kunz T, Kirstein M. A prospective study on ultrasound-guided compression therapy of thrombin injection for treatment of iatrogenic false aneurysms in patients receiving full-dose anti-platelet therapy. *Z Kardiol* 2003; 92(7):564–570
- [51] Görges G, Kunz T. Management of complex, two chamber false aneurysm by thrombin injections. *Heart* 2003 May; 89(5):564. doi:10.1136/heart.89.5.564
- [52] Maleux G, Hendrickx S, Vaninbrouckx J et al. Percutaneous injection of human thrombin to treat iatrogenic femoral pseudoaneurysms: short- and midterm ultrasound follow-up. *Eur Radiol* 2003; 13:209–212
- [53] Cope C, Zeit R. Coagulation of aneurysms by direct percutaneous thrombin injection. *AJR Am J Roentgenol* 1986; 147(2):383–387.
- [53a] Chen DH, Sammel AM, Jain P et al. Cardiologist operated ultrasound guided thrombin injection as a safe and efficacious first line treatment for iatrogenic femoral artery pseudoaneurysms. *Heart Lung Circ* 2015; 24(2):165–72
- [54] Ferguson JD, Whatling PJ, Martin V et al. Ultrasound guided percutaneous thrombin injection of iatrogenic femoral artery pseudoaneurysms after coronary angiography and intervention. *Heart* 2001; 85(4):E5
- [55] Tsetis D, Kochiadakis G, Hamilos M et al. Ultrasound guided percutaneous thrombin injection for the treatment of post-catheterisation femoral pseudoaneurysm. *Hellenic J Cardiol* 2006 Mar-Apr; 47(2):112–3
- [56] Paulson EK, Nelson RC, Mayes CE et al. Sonographically guided thrombin injection of iatrogenic femoral pseudoaneurysms: further experience of a single institution. *AJR Am J Roentgenol* 2001; 177(2):309–316

- [57] Sackett WR, Taylor SM, Coffey CB et al. Ultrasound-guided thrombin injection of iatrogenic femoral pseudoaneurysms: a prospective analysis. *Am Surg* 2000 Oct; 66(10):937–40; Discussion 940–942
- [58] Sheiman RG, Brophy DP. Treatment of iatrogenic femoral pseudoaneurysms with percutaneous thrombin injection: experience in 54 patients. *Radiology* 2001; 219(1):123–127
- [59] Taylor BS, Rhee RY, Muluk S et al. Thrombin injection versus compression of femoral artery pseudoaneurysms. *J Vasc Surg* 1999; 30(6):1052–1059
- [60] Pope M, Johnston KW. Anaphylaxis after thrombin injection of a femoral pseudoaneurysm: recommendations for prevention. *J Vasc Surg* 2000; 32(1):190–191
- [61] Sadiq S, Ibrahim W. Thromboembolism complicating thrombin injection of femoral artery pseudoaneurysm: management with intraarterial thrombolysis. *J Vasc Interv Radiol* 2001; 12(5):633–636
- [62] Tamim WZ, Arbid EJ, Andrews LS et al. Percutaneous induced thrombosis of iatrogenic femoral pseudoaneurysms following catheterization. *Ann Vasc Surg* 2000; 14(3):254–259
- [63] Sievert H, Baser A, Pfeil W et al. The treatment of iatrogenic spurious aneurysm of the femoral artery by direct thrombin injection. *Dtsch Med Wochenschr* 2000; 125(27):822–825
- [64] Ortel TL, Charles LA, Keller FG et al. Topical thrombin and acquired coagulation factor inhibitors: clinical spectrum and laboratory diagnosis. *Am J Hematol* 1994; 45(2):128–135
- [65] Spotnitz WD, Dalton MS, Baker JW et al. Successful use of fibrin glue during 2 years of surgery at a university medical center. *Am Surg* 1989; 55(3):166–168
- [66] Zins M, Vilgrain V, Gayno S et al. US-guided percutaneous liver biopsy with plugging of the needle track: a prospective study in 72 high-risk patients. *Radiology* 1992; 84(3):841–843
- [67] Dorion RP, Hamati HF, Landis B et al. Risk and clinical significance of developing antibodies induced by topical thrombin preparations. *Arch Pathol Lab Med* 1998; 122(10):887–894
- [68] Ortel TL, Moore KD, Quinn-Allen MA et al. Inhibitory anti-factor V antibodies bind to the factor V C2 domain and are associated with hemorrhagic manifestations. *Blood* 1998; 91(11):4188–4196