

13 Angeborene Herzfehler im Erwachsenenalter

- [1] <http://www.kompetenznetz-ahf.de/wir/register> (21.2.2022)
- [2] <https://www.awmf.org/leitlinien/aktuelle-leitlinien/ll-liste/deutsche-gesellschaft-fuer-paediatrische-kardiologie-ev.html>
- [3] Bonhoeffer P, Boudjemline Y, Saliba Z et al. Percutaneous replacement of pulmonary valve in a right-ventricle to pulmonary-artery prosthetic conduit with valve dysfunction. *Lancet* 356 (9239):1403–1405, 2000
- [4] Coats L, Tsang V, Khambadkone S et al. The potential impact of percutaneous pulmonary valve stent implantation on right ventricular outflow tract re-intervention. *Eur J Cardiothorac Surg* 27:536–543, 2005
- [5] Bergersen L, Benson LN, Gillespie MJ et al. Harmony feasibility trial: acute and short-term outcomes with a self-expanding transcatheter pulmonary valve. *JACC Cardiovasc Interv* 10:1763–1773, 2017
- [6] Kim GB. First human cases of new self-expandable percutaneous pulmonary valve implantation using knitted nitinol-wire stent mounted with a tri-leaflet porcine pericardial valve. *J Amer Coll Cardiol* 69:Suppl A-159, 2017
- [7] Vigl M et al. Sexuality and subjective well-being in male patients with congenital heart disease. *Heart* 14:1179–1183, 2009
- [8] Vigl M, Niggemeyer E, Hager A et al. The importance of socio-demographic factors for the quality of life of adults with congenital heart disease. *Quality Life Res* 20:169–177, 2011
- [9] Müller J, Amberger T, Berg A et al. Physical activity in adults with congenital heart disease and associations with functional outcomes. *Heart* 103:1117–1121, 2017
- [10] Bauer UMM, Körten MA, Diller GP et al. Cardiovascular risk factors in adults with congenital heart defects – recognised but not treated? An analysis of the German National Register for Congenital Heart Defects. *Int J Cardiol* 277:79–84, 2019
- [11] Reiner B, Oberhoffer R, Ewert P et al. Quality of life in young people with congenital heart disease is better than expected. *Arch Dis Child* 104:124–128, 2019
- [12] Baumgartner H, Bonhoeffer P, de Groot NM et al. ESC guidelines for the management of grown-up congenital heart disease (new version 2010). *Eur Heart J* 31:2915–2957, 2010
- [13] Bayer O, Loogen F, Wolter HH. Die Herzkatheterisierung bei angeborenen und erworbenen Herzfehlern, 2. Aufl. Thieme, Stuttgart, S. 54–56, 1967
- [14] Morray BH, Jones TK, Coe JY et al. Implantation of the Melody transcatheter pulmonary valve PB1016 in patients with dysfunctional right ventricular outflow tract conduits. *Catheter Cardiovasc Interv* 93:474–480, 2019
- [15] Hascoët S, Dalla Pozza R, Bentham J et al. Early outcomes of percutaneous pulmonary valve implantation using the Edwards SAPIEN 3 transcatheter heart valve system. *EuroIntervention* 14:1378–1385, 2019
- [16] Hascoët S, Karsenty C, Tortigue M et al. A modified procedure for percutaneous pulmonary valve implantation of the Edwards SAPIEN 3 valve. *EuroIntervention* 14:1386–1388, 2019
- [17] Plessis J, Hascoët S, Baruteau A et al. Edwards SAPIEN transcatheter pulmonary valve implantation: results from a french registry. *JACC Cardiovasc Interv* 11:1909–1916, 2018
- [18] Morgan G, Prachasilchai P, Promphan W et al. Medium-term results of percutaneous pulmonary valve implantation using the Venus P-valve™: international experience. *EuroIntervention* 14:1363–1370, 2019
- [19] Kim GB, Song MK, Bae EJ et al. Successful feasibility human trial of a new self-expandable percutaneous pulmonary valve (Pulsta Valve) implantation using knitted

- nitinol wire backbone and trileaflet α -gal-free porcine pericardial valve in the native right ventricular outflow tract. *Circ Cardiovasc Interv* 11:e006494, 2018
- [20] Lababidi Z, Wu JR, Walls TJ. Percutaneous balloon aortic valvuloplasty: results in 23 patients. *Am J Cardiol* 53:194–197, 1984
- [21] Walls JT, Lababidi Z, Curtes JJ et al. Assessment of percutaneous balloon pulmonary and aortic valvuloplasty. *J Thorac Cardiovasc Surg* 88:352–356, 1984
- [22] Rupprath G, Neuhaus KL. Percutaneous balloon valvuloplasty for aortic valve stenosis in infancy. *Am J Cardiol* 55:1655–1656, 1985
- [23] Vogt A, Rupprath G, Tebbe U et al. Verlaufsbeobachtung nach Dilatation angeborener und erworbener Aortenstenosen. *Dtsch Med Wochenschr* 113:1956–1959, 1988
- [24] Ten Harkel AD, Berkhout M, Hop WC et al. Congenital valvular aortic stenosis: limited progression during childhood. *Arch Dis Child* 94:531–535, 2009
- [25] van der Linde D, Andrinopoulou ER, Oechslin EN et al. Congenital valvular aortic stenosis in young adults: predictors for rate of progression of stenosis and aortic dilatation. *Int J Cardiol* 168:863–870, 2013
- [26] Sen S, Garg S, Rao SG et al. Native aortic coarctation in neonates and infants: immediate and midterm outcomes with balloon angioplasty and surgery. *Ann Pediatr Cardiol* 11:261–266, 2018
- [27] Bugeja J, Cutajar D, Zahra C et al. Aortic stenting for neonatal coarctation of the aorta – when should this be considered? *Images Paediatr Cardiol* 18:1–4, 2016
- [28] Ewert P, Berger F, Kretschmar O et al. Stentimplantation als Therapie der ersten Wahl bei Erwachsenen mit Aortenisthmusstenose? *Z Kardiol* 92:48–52, 2003
- [29] Haas NA, Lewin MA, Knirsch W et al. Initial experience using the NuMED Cheat-ham Platinum (CP) stent for interventional treatment of coarctation of the aorta in children and adolescents. *Z Kardiol* 94:113–120, 2005
- [30] Salcher M, Naci H, Law TJ et al., Cardio-proof Consortium. Balloon dilatation and stenting for aortic coarctation: a systematic review and meta-analysis. *Circ Cardiovasc Interv* 9:e003153, 2016
- [31] Morray HB. Ventricular septal defect closure devices, techniques, and outcomes. *Intervent Cardiol Clin* 8:1–10, 2019
- [32] Saurav A, Kaushik M, Mahesh Alla V et al. Comparison of percutaneous device closure versus surgical closure of peri-membranous ventricular septal defects: A systematic review and meta-analysis. *Catheter Cardiovasc Interv* 86:1048–1056, 2015
- [33] Gray RG, Menon SC, Johnson JT et al. Acute and midterm results following percutaneous device closure of muscular ventricular septal defects: a multicenter PICES investigation. *Catheter Cardiovasc Interv* 90:281–289, 2017
- [34] Agrawal H, Waller BR 3rd, Surendan S et al. New patent ductus arteriosus closure devices and techniques. *Interv Cardiol Clin* 8:23–32, 2019
- [35] Cesnjevar R, Dietrich S. S2k-Leitlinie: Persistierender Ductus arteriosus im Kindes- und Jugendalter. *Bayer Ärztzbl* 640–645, 2016
- [36] Fontan F, Baudet E. Surgical repair of tricuspid atresia. *Thorax* 6:240–249, 1971
- [36a] Rychik J, Atz AM, Celermajer DS et al., American Heart Association Council on Cardiovascular Disease in the Young and Council on Cardiovascular and Stroke Nursing. Evaluation and management of the child and adult with fontan circulation: a scientific statement from the American Heart Association. *Circulation* 140:e234–e284, 2019
- [36b] Freedom RM, Nykanen D, Benson LN. The physiology of the bidirectional cavopulmonary connection. *Ann Thorac Surg* 66:664–667, 1998
- [37] Taussig BH, Bing RJ. Complete transposition of the aorta and a levoposition of the pulmonary artery – clinical, physiological, and pathological findings. *Am Heart J* 37:551–559, 1949
- [38] Gaspardone A, De Marco F, Sgueglia GA et al. Novel percutaneous suture-mediated patent foramen ovale closure technique: early results of the NobleStitch EL Italian Registry. *EuroIntervention* 14:e272–279, 2018
- [39] Welch TR, Nugent AW, Veeram Reddy SR. Biodegradable stents for congenital heart disease. *Interv Cardiol Clin* 8:81–94, 2019
- [40] Sigler M, Söderberg B, Schmitt B et al. Carag bioresorbable septal occluder (CBSO):

- histopathology of experimental implants. *EuroIntervention* 13:1655–1661, 2018
- [41] Germans T, Wilde AA, Dijkmans PA et al. Structural abnormalities of the inferoseptal left ventricular wall detected by cardiac magnetic resonance imaging in carriers of hypertrophic cardiomyopathy mutations. *J Am Coll Cardiol* 48:2518–2523, 2006
- [42] Johansson B, Maceira AM, Babu-Narayan SV et al. Clefts can be seen in the basal inferior wall of the left ventricle and the interventricular septum in healthy volunteers as well as patients by cardiovascular magnetic resonance. *J Am Coll Cardiol* 50:1294–1295, 2007
- [43] Aquaro GD, Di Bella G, Strata E. Cardiac magnetic resonance findings in isolated congenital left ventricular diverticuli. *Int J Cardiovasc Imaging* 23:43–47, 2007
- [44] Basso C, Marra MP, Thiene G. Myocardial clefts, crypts, or crevices: once again, you see only what you look for. *Circ Cardiovasc Imaging* 7:217–219, 2014
- [45] Teare D. Asymmetrical hypertrophy of the heart in young adults. *Br Heart J* 20:1–8, 1958
- [46] Wein M, Wolf-Puetz A, Niehues R et al. Multiple linksventrikuläre inferoseptale Clefts. *Herz* 36:438–443, 2011
- [47] Seraphim A, Papachristidis A, Bratt N et al. Significance of left ventricular clefts: a case report. *J Cardiol Cases* 9:138–142, 2014
- [48] Petryka J, Baksi AJ, Prasad SK et al. Prevalence of inferobasal myocardial crypts among patients referred for cardiovascular magnetic resonance. *Circ Cardiovasc Imaging* 7:259–264, 2014
- [49] Maron MS, Rowin EJ, Lin D et al. Prevalence and clinical profile of myocardial crypts in hypertrophic cardiomyopathy. *Circ Cardiovasc Imaging* 5:441–447, 2012
- [50] Child N, Muhr T, Sammut E et al. Prevalence of myocardial crypts in a large retrospective cohort study by cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 16:66, 2014
- [51] Deykina D, Merzhina EA, Arkhipova I et al. Occurrence and significance of myocardial clefts detected on coronary CTA. *ECR* 2012. doi: 10.1594/ecr2012/C-2232, 2012
- [52] Brouwer WP, Germans T, Head MC et al. Multiple myocardial crypts on modified long-axis view are a specific finding in pre-hypertrophic HCM mutation carriers. *Eur Heart J Cardiovasc Imaging* 13:292–297, 2012
- [53] Jenni R, Oechslin E, Schneider J et al. Echocardiographic and pathoanatomical characteristics of isolated left ventricular non-compaction: a step towards classification as a distinct cardiomyopathy. *Heart* 86:666–671, 2001
- [54] Petersen SE, Selvanayagam JB, Wiesmann F et al. Left ventricular non-compaction: insights from cardiovascular magnetic resonance imaging. *J Am Coll Cardiol* 46:101–105, 2005
- [55] Srichai MB, Hecht EM, Kim DC et al. Ventricular diverticula on cardiac CT: more common than previously thought. *AJR Am J Roentgenol* 189:204–208, 2007
- [56] Ohlow MA. Congenital left ventricular aneurysms and diverticula: definition, pathophysiology, clinical relevance and treatment. *Cardiology* 106:63–72, 2006
- [57] Ohlow MA, Secknus MA, Geller JC et al. Prevalence and outcome of congenital left ventricular aneurysms and diverticula in an adult population. *Cardiology* 112:287–293, 2009
- [58] Krasemann T, Gehrmann J, Fenge H et al. Ventricular aneurysm or diverticulum? Clinical differential diagnosis. *Pediatr Cardiol* 22:409–411, 2001
- [59] Grant RT. An unusual anomaly of the coronary vessels in the malformed heart for a child. *Heart* 13:273–283, 1926
- [60] Engberding R, Bender F. Identification of a rare congenital anomaly of the myocardium by two-dimensional echocardiography: persistence of isolated myocardial sinusoids. *Am J Cardiol* 53:1733–1734, 1984
- [61] Engberding R, Stöllberger C, Ong P et al. Isolierte Noncompaction-Kardiomyopathie. *Dtsch Arztebl Int* 107:206–213, 2010
- [62] Chin TK, Perloff JK, Williams RG et al. Isolated noncompaction of left ventricular myocardium: a study of eight cases. *Circulation* 82:507–513, 1990
- [63] Dusek J, Ostádal B, Duskova M. Postnatal persistence of spongy myocardium with embryonic blood supply. *Arch Pathol* 99:312–317, 1975
- [64] Sedmera D, McQuinn T. Embryogenesis of the heart muscle. *Heart Fail Clin* 4:235–245, 2008

- [65] Shariati J, Schlosser T, Erbel R. „Non-compaction“-Kardiomyopathie. Herz 40:583–590, 2015
- [66] Murphy RT, Thaman R, Blanes JG et al. Natural history and familial characteristics of isolated left ventricular non-compaction. Eur Heart J 26:187–192, 2005
- [67] Towbin JA, Lorts A, Jefferies JL. Left ventricular non-compaction cardiomyopathy. Lancet 386(9995):813–825, 2015
- [68] Carrilho-Ferreira P, Almeida AG, Pinto FJ. Non-compaction cardiomyopathy: prevalence, prognosis, pathoetiology, genetics, and risk of cardioembolism. Curr Heart Fail Rep 11:393–403, 2014
- [69] Jenni R, Oechslin E, Schneider J et al. Echocardiographic and pathoanatomical characteristics of isolated left ventricular non-compaction: a step towards classification as a distinct cardiomyopathy. Heart 86:666–671, 2001
- [70] Stöllberger C, Finsterer J. Left ventricular hypertrabeculation/noncompaction. J Am Soc Echocardiogr 17:91–100, 2004
- [71] Petersen SE, Selvanayagam JB, Wiesmann F et al. Left ventricular non-compaction: insights from cardiovascular magnetic resonance imaging. J Am Coll Cardiol 46:101–105, 2005
- [72] Jacquier A, Thuny F, Jop B et al. Measurement of trabeculated left ventricular mass using cardiac magnetic resonance imaging in the diagnosis of left ventricular non-compaction. Eur Heart J 31:1098–1104, 2010
- [73] Stacey RB, Andersen MM, St Clair M et al. Comparison of systolic and diastolic criteria for isolated LV noncompaction in CMR. JACC Cardiovasc Imaging 6:931–940, 2013
- [73a] Weir-McCall JR, Yeap PM, Papagiorcopulo C et al. Left ventricular noncompaction: anatomical phenotype or distinct cardiomyopathy? J Am Coll Cardiol 68:2157–2165, 2016
- [74] Captur G, Muthurangu V, Cook C et al. Quantification of left ventricular trabeculae using fractal analysis. J Cardiovasc Magn Reson 15:36, 2013
- [75] Kawel-Boehm N, McClelland RL, Zemrak F et al. Hypertrabeculated left ventricular myocardium in relationship to myocardial function and fibrosis: The Multi-Ethnic Study of Atherosclerosis. Radiology 284:667–675, 2017
- [76] Ivanov A, Dabiesingh DS, Bhumireddy GP et al. Prevalence and prognostic significance of left ventricular noncompaction in patients referred for cardiac magnetic resonance imaging. Circ Cardiovasc Imaging 10(9):e006174, 2017
- [77] Oechslin EN, Attenhofer Jost CH, Rojas JR et al. Long-term follow-up of 34 adults with isolated left ventricular noncompaction: a distinct cardiomyopathy with poor prognosis. J Am Coll Cardiol 36:493–500, 2000
- [78] Mohrs OK, Magedanz A, Schlosser T. Non-compaction of the left ventricular myocardium detected by 64-slice multidetector computed tomography. Clin Cardiol 30:48, 2007
- [79] Melendez-Ramirez G, Castillo-Castellon F, Espinola-Zavaleta N et al. Left ventricular noncompaction: a proposal of new diagnostic criteria by multidetector computed tomography. J Cardiovasc Comput Tomogr 6:346–354, 2012
- [80] Engberding R, Yelbuz TM, Breithardt G. Isolated noncompaction of the left ventricular myocardium – a review of the literature two decades after the initial case description. Clin Res Cardiol 96:481–488, 2007
- [81] Steffel J, Duru F. Rhythm disorders in isolated left ventricular noncompaction. Ann Med 44:101–108, 2012
- [82] Onay OS, Yildirim I, Beken B et al. Successful implantation of an intracardiac defibrillator in an infant with long QT syndrome and isolated noncompaction of the ventricular myocardium. Pediatr Cardiol 34:189–193, 2013
- [83] Jefferies JL. Are we getting closer to risk stratification in left ventricular noncompaction cardiomyopathy? J Am Heart Assoc 7:e010608, 2018
- [84] Zemrak F, Ahlman MA, Captur G et al. The relationship of left ventricular trabeculation to ventricular function and structure over a 9.5-year follow-up: the MESA study. J Am Coll Cardiol 64:1971–1980, 2014
- [85] Thavendiranathan P, Dahiya A, Phelan D et al. Isolated left ventricular non-compaction controversies in diagnostic criteria, adverse outcomes and management. Heart 99:681–689, 2013
- [86] Columbus MR. De re anatomica. XV, 265, 1559

- [87] Auch-Schwelk W, Bonzel T, Krause T et al. Differential diagnosis of chest pain and diagnostic findings in pericardial defects combined with coronary artery disease. *Clin Cardiol* 11:650–657, 1988
- [87a] Johanssen L. Diseases of the pericardium. A. Development of the pericardial cavity. In: Derra E (Ed), *Handbuch der Thoraxchirurgie*, 614–615. Springer, Berlin, Göttingen, Heidelberg, 1959
- [88] Jaglan A, Leybushkis B, Tajik AJ. Topsy-turvy heart: a case of congenital absence of the pericardium. *Eur Heart J CV Imag* 21:269, 2020
- [89] Van Son JA, Danielson GK, Schaff HV et al. Congenital partial and complete absence of the pericardium. *Mayo Clin Proc* 68:743–747, 1993
- [90] Ellis K, Leeds NE, Himmelstein A. Congenital deficiencies in the parietal pericardium: a review with 2 new cases including successful diagnosis by plain roentgenography. *Am J Roentgenol Radium Ther Nucl Med* 82:125–137, 1959
- [91] Glover LB, Barcia A, Reeves TJ. Congenital absence of the pericardium. A review of the literature with demonstration of a previously unreported fluoroscopic finding. *Am J Roentgenol Radium Ther Nucl Med* 106:542–549, 1969
- [92] Palau P, Domínguez E, García-González P et al. Isolated partial congenital absence of the pericardium: a familial presentation. *Can J Cardiol* 32:1039.e1–2, 2016
- [93] Persigehl M, Erbel R. Die angeborene linksseitige Perikardaplasie. *Fortschr Röntgenstr* 135:541–547, 1981
- [93] Üzünlü I, Büyük Y, Pakiş I et al. Sudden death due to congenital pericardial defect: an autopsy case. *Am J Forensic Med Pathol* 29: 242–244, 2008
- [94] Foo JS, Koh CH, Sahlén A et al. Congenital partial absence of pericardium: a mimic of arrhythmogenic right ventricular cardiomyopathy. *Case Rep Med* 2018:4297280, 2018
- [95] Shah AB, Kronzon I. Congenital defects of the pericardium: a review. *Eur Heart J Cardiovasc Imaging* 16:821–827, 2015
- [96] Huang J, Slaughter MS. Congenital partial absence of the pericardium. *Ann Thorac Surg* 91:e28, 2011
- [97] Ren G, Juan YH, Fei H et al. Congenital absence of the pericardium and tricuspid regurgitation. *QJM* 111:895–897, 2018
- [98] von Bernuth G, Nissen H, Lang D et al. Angeborene Defekte des linksseitigen Perikards. *Z Kardiol* 65:1022–1032, 1976
- [99] Morgan JR, Rogers AK, Forker AD. Congenital absence of the left pericardium: clinical findings. *Ann Intern Med* 74:370–376, 1971
- [100] Han J, Xiang H, Ridley WE et al. Snoopy sign: congenital absence of the left pericardium. *J Med Imaging Radiat Oncol* 62(Suppl 1):47, 2018
- [101] Kim HJ, Cho YS, Cho GY et al. Congenital absence of the pericardium. *J Cardiovasc Ultrasound* 22:36–39, 2014
- [102] Jurko A Jr, Minarik M, Cisarikova V et al. Congenital complete and partial absence of the left pericardium. *Wien Med Wochenschr* 163:426–428, 2013
- [103] Marta L, Ferreira AM, Santos KR. Congenital absence of pericardium: a nomad heart. *Cardiol Young* 25:1415–1417, 2015
- [104] Gatzoulis MA, Munk MD, Merchant N et al. Isolated congenital absence of the pericardium: clinical presentation, diagnosis, and management. *Ann Thorac Surg* 69:1209–1215, 2000
- [105] Abbas AE, Appleton CP, Liu PT et al. Congenital absence of the pericardium: case presentation and review of literature. *Int J Cardiol* 98:21–25, 2005
- [106] Huang J, Slaughter MS. Congenital partial absence of the pericardium. *Ann Thorac Surg* 91:e28, 2011
- [107] De Brabandere K, Czaplaj J, Callens E et al. Coronary artery bypass surgery in a patient with congenital absence of the left pericardium. *Eur J Cardiothorac Surg* 41:e65, 2012
- [108] Tăbăran AF, Nagy AL, Cătoi C et al. Congenital diaphragmatic hernia with concurrent aplasia of the pericardium in a foal. *BMC Vet Res* 11:309, 2015