

17 Behandlungsstrategien in besonderen Fällen

17.1 Akuter Myokardinfarkt

- [1] Roffi M, Patrono C, Collet JP et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J* 37:267–315, 2016
- [2] Thygesen K, Alpert JS, Jaffe AS et al., Executive Group on behalf of the Joint European Society of Cardiology (ESC)/ American College of Cardiology (ACC)/ American Heart Association (AHA)/World Heart Federation (WHF). Task Force for the Universal Definition of Myocardial Infarction. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol* 72:2231–64, 2018
- [3] Goyal A, Gluckman TJ, Tcheng JE. What's in a name? The new ICD-10 (10th Revision of the International Statistical Classification of Diseases and Related Health Problems) codes and type 2 myocardial infarction. *Circulation* 136:1180–2, 2017
- [4] Giannitsis E, Katus H. Troponins: established and novel indications in the management of cardiovascular disease. *Heart* 104:1714–22, 2018
- [5] Ge J, Erbel R, Zamorano J et al. Coronary artery remodeling in atherosclerotic disease: an intravascular ultrasonic study in vivo. *Coron Artery Dis* 4:981–6, 1993
- [6] Hartmann M, von Birgelen C, Mintz GS et al. Relation between baseline plaque burden and subsequent remodelling of atherosclerotic left main coronary arteries: a serial intravascular ultrasound study with long-term (> or = 12 months) follow-up. *Eur Heart J* 27:1778–84, 2006
- [7] Erbel R, Ge J, Görges G et al. Intravascular ultrasound classification of atherosclerotic lesions according to American Heart Association recommendation. *Coron Artery Dis* 10:489–99, 1999
- [8] Ge J, Liu F, Görges G et al. Angiographically „silent“ plaque in the left main coronary artery detected by intravascular ultrasound. *Coron Artery Dis* 6:805–10, 1995
- [9] Virmani R, Kolodgie FD, Burke AP, Farb A, Schwartz SM. Lessons from sudden coronary death: a comprehensive morphological classification scheme for atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* 20:1262–75, 2000
- [10] Erbel R. Vulnerable plaque – vulnerable vessel – vulnerable patient. Plaques prone to rupture in atheroma/fibroatheroma. *Herz* 28:483–7, 2003
- [11] Erbel R, Heusch G. Coronary microembolization – its role in acute coronary syndromes and interventions. *Herz* 24:558–75, 1999
- [12] Erbel R, Heusch G. Coronary microembolization. *J Am Coll Cardiol* 36:22–4, 2000
- [13] Erbel R. Progression and regression of atherosclerotic plaques. New results based on intracoronary ultrasound. *Herz* 40:855–62, 2015
- [14] Lyon AR, Bossone E, Schneider B et al. Current state of knowledge on Takotsubo syndrome: a Position Statement from the Taskforce on Takotsubo Syndrome of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 18:8–27, 2016
- [15] Hayes SN, Kim ESH, Saw J et al. Spontaneous coronary artery dissection: current state of the science: a scientific statement from the American Heart Association. *Circulation* 137:e523–57, 2018
- [16] Hawranek M, Wróbel M, Nożyński J et al. Hemorrhagic myocardial infarction: mortality compared with STEMI patients treated with percutaneous coronary intervention. *J Am Coll Cardiol* 68:426–7, 2016

- [17] Baars T, Möhlenkamp S, Eggebrecht H et al. Diagnostik von Thoraxschmerzen, Teil 2 – Kardiale Ursachen müssen immer als Erstes abgeklärt werden. *Pneumo-News* 5:39–43, 2008
- [18] Möhlenkamp S, Schmerund A, Lehmann S et al. Subclinical coronary atherosclerosis and resting ECG abnormalities in an unselected general population. *Atherosclerosis* 196:786–94, 2008
- [19] Mehilli J, Hamm CW, Landmesser U, Massberg S. Kommentar zu den 2015-Leitlinien der Europäischen Gesellschaft für Kardiologie (ESC) zum Management des akuten Koronarsyndroms ohne ST-Hebungen (NSTEMI-ACS). *Kardiologie* 10:351–8, 2016
- [20] Holmes DR Jr, Berger PB. Troponins, necroptosis, enzyme leaks, creatinine phosphokinase bumps, and infarctlets: what's behind this new lexicon and what does it add? *Circulation* 104:627–9, 2001
- [21] Eggebrecht H, Plicht B, Buck T et al. Echokardiographische Abklärung des Patienten mit akutem Thoraxschmerz auf der Notfallstation. *Intensivmed* 3:64–77, 2006
- [22] Breuckmann F, Post F, Giannitsis E et al. Kriterien der Deutschen Gesellschaft für Kardiologie – Herz- und Kreislaufforschung für „Chest-Pain-Units“. *Kardiologie* 2:389–94, 2008
- [22a] Giannitsis E, Post F, Haerer W et al. Kriterien der Deutschen Gesellschaft für Kardiologie – Herz- und Kreislaufforschung für „Chest Pain Units“. *Kardiologie* 14:446–79, 2020
- [23] Post F, Gori T, Giannitsis E et al. Criteria of the German Society of Cardiology for the establishment of chest pain units: update 2014. *Clin Res Cardiol* 104:918–28, 2015
- [23a] Breuckmann F, Rassaf T, Hochadel M et al. German chest pain unit registry: data review after the first decade of certification. *Herz* 46(Suppl 1):24–32, 2021
- [24] Breuckmann F, Hochadel M, Voigtländer T et al. The use of echocardiography in certified chest pain units: results from the German Chest Pain Unit Registry. *Cardiology* 134:75–83, 2016
- [25] Mehilli J, Hamm CW, Landmesser U, Massberg S. Kommentar zu den 2015-Leitlinien der Europäischen Gesellschaft für Kardiologie (ESC) zum Management des akuten Koronarsyndroms ohne ST-Hebungen (NSTEMI-ACS). *Kardiologie* 10:351–358, 2016
- [25a] Thiele H, Bauersachs J, Mehilli J et al. Kommentar zu den 2020er Leitlinien der Europäischen Gesellschaft für Kardiologie (ESC) zum Management des akuten Koronarsyndroms bei Patienten ohne persistierende ST-Strecken-Hebung. *Kardiologie* 15:19–31, 2021
- [26] Fox KA, Fitzgerald G, Puymirat E et al. Should patients with acute coronary disease be stratified for management according to their risk? Derivation, external validation and outcomes using the updated GRACE risk score. *BMJ Open* 4:e004425, 2014
- [27] http://www.outcomes-umassmed.org/grace/files/GRACE_RiskModel_Coefficients.pdf
- [27a] Collet JP, Thiele H, Barbato E., ESC Scientific Document Group. 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 42:1289–367, 2021. Erratum in: *Eur Heart J* 42:1908, 2021. Erratum in: *Eur Heart J* 42:1925, 2021. Erratum in: *Eur Heart J* 42:2298, 2021
- [28] Subherwal S, Bach RG, Chen AY et al. Baseline risk of major bleeding in non-ST-segment-elevation myocardial infarction: the CRUSADE (Can Rapid risk stratification of Unstable angina patients Suppress ADverse outcomes with Early implementation of the ACC/AHA guidelines) bleeding score. *Circulation* 119:1873–82, 2009
- [29] <https://www.mdcalc.com/calc/1784/crusade-score-post-mi-bleeding-risk>
- [30] Al-Daydamony MM, Farag El-Sayed M. CRUSADE bleeding score as a predictor of bleeding events in patients with acute coronary syndrome in Zagazig University Hospital. *Indian Heart Journal* 68:632–8, 2016
- [31] Löwel H, Meisinger C. Epidemiologie und demographische Entwicklung am Beispiel kardiovaskulärer Erkrankungen in Deutschland. *Med Klin (Munich)* 101:804–11, 2006
- [31a] Andresen D. 32. Herzbericht 2020. Thieme, Stuttgart, 2021
- [32] Dudas K, Lappas G, Stewart S et al. Trends in out-of-hospital deaths due to coronary heart disease in Sweden (1991 to 2006). *Circulation* 123:46–52, 2011

- [33] Zeymer U, Vogt A, Zahn R et al. Predictors of in-hospital mortality in 1.333 patients with acute myocardial infarction complicated by cardiogenic shock treated with primary percutaneous coronary intervention (PCI). Results of the primary PCI registry of the Arbeitsgemeinschaft Leitende Kardiologische Krankenhausärzte (ALKK). *Eur Heart J* 25:322–8, 2004
- [34] Liosis S, Bauer T, Schiele R et al. Predictors of 1-year mortality in patients with contemporary guideline-adherent therapy after acute myocardial infarction: results from the OMEGA study. *Clin Res Cardiol* 102:671–7, 2013
- [35] Jacksch R, Naber CK, Koslowski B et al. Akutintervention (Primär-PCI) im Rahmen des Essener Herzinfarktverbundes. *Herz* 33:110–4, 2008
- [36] Hailer B, Naber C, Koslowski B et al. Gender-related differences in patients with ST-elevation myocardial infarction: results from the registry study of the ST elevation myocardial infarction network Essen. *Clin Cardiol* 34:294–301, 2011
- [37] Scholz KH, Maier SKG, Maier LS et al. Impact of treatment delay on mortality in ST-segment elevation myocardial infarction (STEMI) patients presenting with and without haemodynamic instability: results from the German prospective, multicentre FITT-STEMI trial. *Eur Heart J* 39:1065–74, 2018
- [38] Byrne RA, Rossello X, Coughlan JJ et al. ESC Scientific Document Group. 2023 ESC Guidelines for the management of acute coronary syndromes. *Eur Heart J* 44:3720–826, 2023
- [39] Kelm M, Bauersachs J, Kastrati A et al. 2017 ESC Pocket Guidelines. Therapie des akuten Herzinfarktes bei Patienten mit persistierender ST-Streckenhebung. Version 2017. Kurzfassung der 2017 ESC Guidelines for the management of acute myocardial infarction in patients with ST-segment elevation. (European Heart Journal 2017). Börm Bruckmeier Verlag, Grünwald, 2016
- [40] Kelm M, Kastrati A, Nef H et al. für Kommission für Klinische Kardiologische Medizin. Kommentar zu den Leitlinien 2017 der Europäischen Gesellschaft für Kardiologie (ESC) zur Therapie des akuten Herzinfarktes bei Patienten mit ST-Streckenhebung. *Kardiologie* 12:145–9, 2018
- [41] Thiele H, Akin I, Sandri M et al. PCI Strategies in patients with acute myocardial infarction and cardiogenic shock. *N Engl J Med* 377:2419–32, 2017
- [42] Beltrane JF. Assessing patients with myocardial infarction and nonobstructive coronary arteries (MINOCA). *J Intern Med* 273:182–5, 2013
- [43] Pasupathy S, Air T, Dreyer RP et al. Systematic review of patients presenting with suspected myocardial infarction and non-obstructive coronary arteries. *Circulation* 131:861–70, 2015
- [44] Pasupathy S, Air T, Dreyer RP et al. Response to letter regarding article, „Systematic review of patients presenting with suspected myocardial infarction and non-obstructive coronary arteries“. *Circulation* 132:e232, 2015
- [45] Agewall S, Beltrane JF, Reynolds HR et al. ESC working group position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J* 38:143–53, 2017
- [46] Breuckmann F, Nassenstein K, Bucher C et al. Systematic analysis of functional and structural changes after coronary microembolization: a cardiac magnetic resonance imaging study. *JACC Cardiovasc Imaging* 2:121–30, 2009
- [47] Collste O, Sörensson P, Frick M et al. Myocardial infarction with normal coronary arteries is common and associated with normal findings on cardiovascular magnetic resonance imaging: results from the Stockholm Myocardial Infarction with Normal Coronaries study. *J Intern Med* 273:189–96, 2013
- [48] Agewall S, Giannitsis E, Jernberg T et al. Troponin elevation in coronary vs. non-coronary disease. *Eur Heart J* 32:404–11, 2011
- [49] Ouldzein H, Elbaz M, Roncalli J et al. Plaque rupture and morphological characteristics of the culprit lesion in acute coronary syndromes without significant angiographic lesion: analysis by intravascular ultrasound. *Ann Cardiol Angeiol (Paris)* 61:20–6, 2012
- [50] Reynolds HR, Srichai MB, Iqbal SN et al. Mechanisms of myocardial infarction in women without angiographically obstructive coronary artery disease. *Circulation* 124:1414–25, 2011

- [51] Davies MJ, Thomas AC. Plaque fissuring – the cause of acute myocardial infarction, sudden ischaemic death, and crescendo angina. *Br Heart J* 53:363–73, 1985
- [52] Davies MJ, Thomas AC, Knapman PA et al. Intramyocardial platelet aggregation in patients with unstable angina suffering sudden ischemic cardiac death. *Circulation* 73:418–27, 1986
- [53] Davies MJ, Bland JM, Hangartner JR et al. Factors influencing the presence or absence of acute coronary artery thrombi in sudden ischaemic death. *Eur Heart J* 10:203–8, 1989
- [54] Falk E, Shah PK, Fuster V. Coronary plaque disruption. *Circulation* 92:657–71, 1995
- [55] Davies MJ. The pathological basis of angina pectoris. *Cardiovasc Drugs Ther* 3(Suppl 1):249–55, 1989
- [56] Ge J, Chirillo F, Schwedtmann J et al. Screening of ruptured plaques in patients with coronary artery disease by intravascular ultrasound. *Heart* 81:621–7, 1999
- [57] Erbel R, Ge J, Gorge G et al. Intravascular ultrasound classification of atherosclerotic lesions according to American Heart Association recommendation. *Coron Artery Dis* 1:489–99, 1999
- [58] Erbel R. Vulnerable Plaque – vulnerables Gefäß – vulnerabler Patient. *Herz* 28:483–7, 2003
- [59] Isner JM, Kishel J, Kent KM et al. Accuracy of angiographic determination of left main coronary arterial narrowing. Angiographic – histologic correlative analysis in 28 patients. *Circulation* 63:1056–64, 1981
- [60] Opolski MP, Spiwak M, Marczak M et al. Mechanisms of myocardial infarction in patients with nonobstructive coronary artery disease: results from the optical coherence tomography study. *ACC Cardiovasc Imaging* 12:2210–21, 2019
- [61] Pathik B, Raman B, Mohd Amin NH et al. Troponin-positive chest pain with unobstructed coronary arteries: incremental diagnostic value of cardiovascular magnetic resonance imaging. *Eur Heart J Cardiovasc Imaging* 17:1146–52, 2016
- [62] Dastidar AG, Baritussio A, De Garate E et al. Prognostic role of CMR and conventional risk factors in myocardial infarction with nonobstructed coronary arteries. *JACC Cardiovasc Imaging* 12:1973–82, 2017
- [63] Roffi M, Patrono C, Collet JP et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: task force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 37:267–315, 2016
- [64] Nicoli G, Scalone G, Crea F. Acute myocardial infarction with no obstructive coronary atherosclerosis: mechanism and management. *Eur Heart J* 36:475–81, 2015
- [64a] Lascarrou JB, Merdji H, Le Gouge A et al., CRICS-TRIGGERSEP Group. Targeted temperature management for cardiac arrest with nonshockable rhythm. *N Engl J Med* 381:2327–37, 2019
- [65] Corpus RA, House JA, Marso SP et al. Multivessel percutaneous coronary intervention in patients with multivessel disease and acute myocardial infarction. *Am Heart J* 48:493–500, 2004
- [66] Stone GW, Grines CL, Cox DA et al. Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med* 46:957–66, 2002
- [67] Glover GW, Thomas RM, Vamvakas G. Intravascular versus surface cooling for targeted temperature management after out-of-hospital cardiac arrest - an analysis of the TTM trial data. *Crit Care* 20:381, 2016
- [68] Arrich J, Holzer M, Havel C et al. Pre-hospital versus in-hospital initiation of cooling for survival and neuroprotection after out-of-hospital cardiac arrest. *Cochrane Database Syst Rev* 3:CD010570, 2016
- [69] Kim KH, Shin SD, Song KJ et al. Cooling methods of targeted temperature management and neurological recovery after out-of-hospital cardiac arrest: A nationwide multicenter multi-level analysis. *Resuscitation* 125:56–65, 2018
- [70] Lindsay PJ, Buell D, Scales DC. The efficacy and safety of pre-hospital cooling after out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Crit Care* 22:66, 2018
- [71] Nolan JP, Soar J, Cariou A et al., European Resuscitation Council and European Society of Intensive Care Medicine Guidelines for Post-resuscitation Care 2015: Section 5 of the European Resuscitation

- Council Guidelines for Resuscitation 2015. Resuscitation 95:202–22, 2015
- [72] Drinhaus H, Stratogianni A, Schick V et al. Postreanimationstherapie: Klare neue Strategien, aber noch offene Fragen. Dtsch Arztebl 113:14, 2016
- [73] Akers WS, Oh JJ, Oestreich JH et al. Pharmacokinetics and pharmacodynamics of a bolus and infusion of cangrelor: a direct, parenteral P2Y₁₂ receptor antagonist. J Clin Pharmacol 50:27–35, 2010
- [74] Droppa M, Geisler T. Cangrelor – Indikationen 2019. J f Kardiologie 26:307–310, 2019
- [75] Topol EJ, Neumann FJ, Montalescot G. A preferred reperfusion strategy for acute myocardial infarction. J Am Coll Cardiol 2:1886–9, 2003
- [76] Vaduganathan M, Qamar A, Singh A et al. Cangrelor Use Since FDA Approval: A Single-Center, Real-World Experience at a Tertiary Care Hospital. J Am Coll Cardiol 69:463–464, 2017
- [77] Burzotta F, Crea F. Thrombus-aspiration: a victory in war against no reflow. Lancet 371:1889–90, 2008
- [78] Zijlstra F, Gonnings NL. Thrombus Aspiration during Percutaneous coronary intervention in Acute myocardial infarction Study (TAPAS). Lancet 371:1915–20, 2008
- [79] Mahmoud KD, Jolly SS, James S et al. Clinical impact of direct stenting and interaction with thrombus aspiration in patients with ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention: Thrombectomy Trialists Collaboration. Eur Heart J 39:2472–9, 2018
- [80] Thielmann M, Kottenberg E, Kleinbongard P et al. Cardioprotective and prognostic effects of remote ischaemic preconditioning in patients undergoing coronary artery bypass surgery: a single-centre randomised, double-blind, controlled trial. Lancet 382:597–604, 2013
- [81] Kholmukhamedov A, Logdon C, Hu J et al. Cyclosporin A in left ventricular remodeling after myocardial infarction. Am J Physiol Heart Circ Physiol 306:H53–9, 2014
- [82] Cung TT, Morel O, Cayla G et al. Cyclosporine before PCI in patients with acute myocardial infarction. N Engl J Med 373:1021–31, 2015
- [83] Yingzhong C, Lin C, Chunbin W. Clinical effects of cyclosporine A on reperfusion injury in myocardial infarction: a meta-analysis of randomized controlled trials. Springerplus 5:1117, 2016

17.2 No-Reflow-Phänomen

- [1] Jaffe R, Charron T, Puley G et al. Microvascular obstruction and the „no-reflow“ phenomenon after percutaneous coronary intervention. Circulation 17:3152–6, 2008
- [2] Lincoff AM, Topol EJ. Illusion of reperfusion. Does anyone achieve optimal reperfusion during acute myocardial infarction? Circulation 8:1361–74, 1993
- [3] Kloner DE, Dai W, Hale SL. No-reflow phenomenon. A new target for therapy of acute myocardial infarction independent of myocardial infarct size. J CV Pharmacol Therapeut 23:273–6, 2018
- [4] Piana RN, Paik GY, Moscucci M et al. Incidence and treatment of „no-reflow“ after percutaneous coronary intervention. Circulation 9:2514–8, 1994
- [5] Mehta RH, Ou FS, Peterson ED et al. Clinical significance of post-procedural TIMI flow in patients with cardiogenic shock undergoing primary percutaneous coronary intervention. JACC Cardiovasc Interv 2:56–64, 2009
- [6] Haeck JD, Koch KT, Bilodeau L et al. Randomized comparison of primary percutaneous coronary intervention with combined proximal embolic protection and thrombus aspiration versus primary percutaneous coronary intervention alone in ST-segment elevation myocardial infarction: the PREPARE (PRoximal Embolic Protection in Acute myocardial infarction and Resolution of ST-Elevation) study. JACC Cardiovasc Interv 2:934–43, 2009
- [7] Halkin A, Stone GW, Grines CL et al. Prognostic implications of creatinine kinase elevation after primary percutaneous coronary intervention for acute myocardial infarction: the CADILLAC risk score. J Am Coll Cardiol 7:951–61, 2006
- [8] Bouleti C, Mewton N, Germain S. The no-reflow phenomenon: state of the art. Arch Cardiovasc Dis 108:661–74, 2015
- [8a] Maznyczka AM, Oldroyd KG, Greenwood JP et al. Comparative significance of invasive measures of microvascular injury in

- acute myocardial infarction. *Circ Cardiovasc Interv* 13:e008505, 2020
- [9] Tanaka A, Kawarabayashi T, Nishibori Y et al. No-reflow phenomenon and lesion morphology in patients with acute myocardial infarction. *Circulation* 105:2148–52, 2002
 - [10] Kotani J, Mintz GS, Castagna MT et al. Usefulness of preprocedural coronary lesion morphology as assessed by intravascular ultrasound in predicting thrombolysis in myocardial infarction frame count after percutaneous coronary intervention in patients with Q-wave acute myocardial infarction. *Am J Cardiol* 1:870–2, 2003
 - [11] Jennings RB, Ganote CE, Reimer KA. Ischemic tissue injury. *Am J Pathol* 1:179–98, 1975
 - [12] Taniyama Y, Ito H, Iwakura K et al. Beneficial effect of intracoronary verapamil on microvascular and myocardial salvage in patients with acute myocardial infarction. *JACC* 30:1193–9, 1997
 - [13] Leinweber K, Böse D, Vogelsang M et al. Intense vasoconstriction in response to aspirate from stented saphenous vein aortocoronary bypass grafts. *J Am Coll Cardiol* 7:981–6, 2006
 - [14] Böse D, von Birgelen C, Zhou XY et al. Impact of atherosclerotic plaque composition on coronary microembolization during percutaneous coronary interventions. *Bas Research Cardiol* 103:587–97, 2008
 - [15] Yu SJ, Buerck JP, O'Rear EA et al. Possible erythrocyte contributions to and exacerbation of the post-thrombolytic no-reflow phenomenon. *Biorheology* 54:81–93, 2018
 - [16] Horn P, Baars T, Kahlert P et al. Release of intracoronary microparticles during stent implantation into stable atherosclerotic lesions under protection with an aspiration device. *PLoS One* 10:e0124904, 2015
 - [17] Caiazzo G, Musci RL, Frediani L et al. State of the Art: No-Reflow Phenomenon. *Cardiol Clin* 38:563–573, 2020
 - [18] Ren F, Mu N, Zhang X et al. Increased platelet-leukocyte aggregates are associated with myocardial no-reflow in patients with ST elevation myocardial infarction. *Am J Med Sci* 352:261–6, 2016
 - [19] Kulickowski W, Miszalski-Jamka K, Kaczmarek J et al. No-reflow and platelet reactivity in diabetic patients with ST-segment elevation myocardial infarction: is there a link? *Postępy Kardiologii Interwencyjnej* 13:326–30, 2017
 - [20] Lee WW, Marinelli B, van der Laan AM et al. PET/MRI of inflammation in myocardial infarction. *J Am Coll Cardiol* 59:153–63, 2012
 - [21] Hawranek M, Wróbel M, Nożyński J et al. Hemorrhagic myocardial infarction: mortality compared with STMI patients treated with percutaneous coronary intervention. *J Am Coll Cardiol* 68:426–7, 2016
 - [22] Abbate A, Kontos MC, Biondi-Zoccai GG. No-reflow: the next challenge in treatment of ST-elevation acute myocardial infarction. *Eur Heart J* 9:1795–7, 2008
 - [23] Limbruno U, De Carlo M, Pistoletti S et al. Distal embolization during primary angioplasty: histopathologic features and predictability. *Am Heart J* 50:102–8, 2005
 - [24] Yip HK, Chen MC, Chang HW et al. Angiographic morphologic features of infarct-related arteries and timely reperfusion in acute myocardial infarction: predictors of slow-flow and no-reflow phenomenon. *Chest* 22:1322–32, 2002
 - [25] Mizote I, Ueda Y, Ohtani T et al. Distal protection improved reperfusion and reduced left ventricular dysfunction in patients with acute myocardial infarction who had angiographically defined ruptured plaque. *Circulation* 12:1001–7, 2005
 - [26] Fukuda D, Tanaka A, Shimada K et al. Predicting angiographic distal embolization following percutaneous coronary intervention in patients with acute myocardial infarction. *Am J Cardiol* 1:403–7, 2003
 - [27] Zhao JL, Yang YJ, Pei WD et al. The effect of statins on the no-reflow phenomenon: an observational study in patients with hyperglycemia before primary angioplasty. *Am J Cardiovasc Drugs* 81–9, 2009
 - [28] Jeong YH, Kim WJ, Park DW et al. Serum B-type natriuretic peptide on admission can predict the „no-reflow“ phenomenon after primary drug-eluting stent implantation for ST-segment elevation myocardial infarction. *Inter J Cardiol* 141:175–81, 2009
 - [29] Niccoli G, Lanza GA, Spaziani C et al. Baseline systemic inflammatory status and no-reflow phenomenon after percutaneous coronary angioplasty for acute myocardial infarction. *Int J Cardiol* 117:306–11, 2007

- [30] Mauri L, Rogers C, Baim DS. Devices for distal protection during percutaneous coronary revascularization. *Circulation* 13:2651–6, 2006
- [31] Iijima R, Shinji H, Ikeda N et al. Comparison of coronary arterial finding by intravascular ultrasound in patients with „transient no-reflow“ versus „reflow“ during percutaneous coronary intervention in acute coronary syndrome. *Am J Cardiol* 7:29–33, 2006
- [32] Soeda T, Higuma T, Abe N et al. Morphological predictors for no reflow phenomenon after primary percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction caused by plaque rupture. *Eur Heart J Cardiovasc Imaging* 18:103–10, 2017
- [33] Kusuyama T, Kataoka T, Iida H et al. Comparison of temporary occlusion and aspiration system versus the conventional method during coronary stenting for acute myocardial infarction. *Am J Cardiol* 4:1041–3, 2004
- [34] Hong YJ, Jeong MH, Choi YH et al. Impact of plaque components on no-reflow phenomenon after stent deployment in patients with acute coronary syndrome: a virtual histology-intravascular ultrasound analysis. *Eur Heart J* 32:2059–66, 2009
- [35] Ding S, Li Z, Ge H et al. Impact of early ST-segment changes on cardiac magnetic resonance-verified intramyocardial haemorrhage and microvascular obstruction in st-elevation myocardial infarction patients. *Medicine* 94:e1438, 2015
- [36] Ndrepepa G, Tiroch K, Fusaro M et al. 5-year prognostic value of no-reflow phenomenon after percutaneous coronary intervention in patients with acute myocardial infarction. *J Am Coll Cardiol* 55:2383–9, 2010
- [37] Nakazawa G, Tanabe K, Onuma Y et al. Efficacy of culprit plaque assessment by 64-slice multidetector computed tomography to predict transient no-reflow phenomenon during percutaneous coronary intervention. *Am Heart J* 155:1150–7, 2008
- [38] Klein LW, Kern MJ, Berger P et al. Society of cardiac angiography and interventions: suggested management of the no-reflow phenomenon in the cardiac catheterization laboratory. *Cath Cardiovasc Interv* 60:194–201, 2003
- [39] Pasceri V, Pristipino C, Pelliccia F et al. Effects of the nitric oxide donor nitroprusside on no-reflow phenomenon during coronary interventions for acute myocardial infarction. *Am J Cardiol* 5:1358–61, 2005
- [40] Michaels AD, Appleby M, Otten MH et al. Pretreatment with intragraft Verapamil prior to percutaneous coronary intervention of saphenous vein graft lesions: results of the randomized, controlled vasodilator prevention on no-reflow (VAPOR) trial. *J Invasive Cardiol* 4:299–302, 2002
- [41] Kaplan BM, Benzuly KH, Kinn JW et al. Treatment of no-reflow in degenerated saphenous vein graft interventions: comparison of intracoronary Verapamil and nitroglycerin. *Cathet Cardiovasc Diagn* 9:113–8, 1996
- [42] Saito T, Hokimoto S, Ishibashi F et al. Pulse infusion thrombolysis (PIT) for large intracoronary thrombus: preventive effect against the „no flow“ phenomenon in revascularization therapy for acute myocardial infarction. *Jpn Circ J* 5:94–8, 2001
- [43] Lim SY, Bae EH, Jeong MH et al. Effect of combined intracoronary adenosine and nicorandil on no-reflow phenomenon during percutaneous coronary intervention. *Circulation* 8:928–32, 2004
- [44] Patel T, Shah S, Gulati R et al. Perforated balloon technique: a simple and handy technique to combat no-reflow phenomenon in coronary system. *Catheter Cardiovasc Interv* 92:890–4, 2018
- [45] Fischell TA, Haller S, Pulukurthy S et al. Nicardipine and adenosine „flush cocktail“ to prevent no-reflow during rotational atherectomy. *Cardiovasc Revasc Med* 224–8, 2008
- [46] Marzilli M, Orsini E, Marraccini P et al. Beneficial effects of intracoronary adenosine as an adjunct to primary angioplasty in acute myocardial infarction. *Circulation* 01:2154–9, 2000
- [47] Movahed MR, Baweja G. Distal administration of very high doses of intracoronary adenosine for the treatment of resistant no-reflow. *Exp Clin Cardiol* 3:141–3, 2008
- [48] Sakuma T, Motoda C, Tokuyama T et al. Exogenous adenosine triphosphate disodium administration during primary percutaneous coronary intervention reduces no-reflow and preserves left ventricular

- function in patients with acute anterior myocardial infarction: a study using myocardial contrast echocardiography. *Int J Cardiol* 140:200–9, 2010
- [48] Heusch G. Adenosine and maximum coronary vasodilation in humans: myth and misconceptions in the assessment of coronary reserve. *Bas Res Cardiol* 105:1–5, 2010
- [49] Kleinbongard P, Boese D, Baars T et al. Vasoconstrictor potential of coronary aspirate from patients undergoing stenting of saphenous vein aortocoronary bypass grafts and its pharmacological attenuation. *Circ Res* 108:344–52, 2011
- [50] Huang D, Qian J, Ge L et al. REstoration of COronary flow in patients with no-reflow after primary coronary interVENTion of acute myocaRdial infarction (RECOVER). *Am Heart J* 164:394–401, 2012
- [51] Freixa X, Bellera N, Ortiz-Perez JT et al. Ischaemic postconditioning revisited: lack of effects on infarct size following primary percutaneous coronary intervention. *Eur Heart J* 33:103–12, 2012
- [52] Høfsten D, Kelbæk H, Helqvist S et al. The Third DANish Study of Optimal Acute Treatment of Patients with ST-segment Elevation Myocardial Infarction: ischemic postconditioning or deferred stent implantation versus conventional primary angioplasty and complete revascularization versus treatment of culprit lesion only. rationale and design of the DANAMI 3 trial program. *Am Heart J* 169:613–21, 2015
- [53] Hahn JY, Song YB, Kim EK et al. Ischemic postconditioning during primary percutaneous coronary intervention: the effects of postconditioning on myocardial reperfusion inpatients with ST-segment elevation myocardial infarction (POST) randomized trial. *Circulation* 128:1889–96, 2013
- [54] Baim DS, Mauri L, Cutlip DC. Drug-eluting stenting for unprotected left main coronary artery disease: are we ready to replace bypass surgery? *J Am Coll Cardiol* 7:878–81, 2006
- [55] Burzotta F, Crea F. Thrombus-aspiration: a victory in the war against no-reflow. *Lancet* 71:1889–90, 2008
- [56] Vlaar PJ, Svilaas T, Horst IC et al. Cardiac death and reinfarction after 1 year in the thrombus aspiration during percutaneous coronary intervention in acute myocardial infarction study (TAPAS): a 1-year follow-up study. *Lancet* 71:1915–20, 2008
- [57] Svilaas T, Vlaar PJ, Horst IC et al. Thrombus aspiration during primary percutaneous coronary intervention. *N Engl J Med* 58:557–67, 2008
- [58] Frobert O, Lagerqvist B, Olivecrona GK et al. Thrombus aspiration during ST-segment elevation myocardial infarction. *N Engl J Med* 369:1587–97, 2013
- [59] Jolly SS, Cairns JA, Yusuf S et al. Randomized trial of primary PCI with or without routine manual thrombectomy. *N Engl J Med* 372:1389–98, 2013
- [60] Byrne RA, Rossello X, Coughlan JJ et al. ESC Scientific Document Group. 2023 ESC Guidelines for the management of acute coronary syndromes. *Eur Heart J* 44:3720–3826, 2023
- [61] Baim DS, Wahr D, George B et al. Randomized trial of a distal embolic protection device during percutaneous intervention of saphenous vein aorto-coronary bypass grafts. *Circulation* 105:1285–90, 2002
- [62] Silber S, Albertsson P, Avilés FF et al. Guidelines for percutaneous coronary interventions. The Task Force for Percutaneous Coronary Interventions of the European Society of Cardiology. *Eur Heart J* 26:804–47, 2005
- [63] Montalescot G, Sechtem U, Achenbach S et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. Web Addenda. *Eur Heart J* 34:2949–3003, 2013. Erratum in: *Eur Heart J* 35:2260–61, 2014
- [64] Brennan JM, Al-Hejily W, Dai D et al. Three-year outcomes associated with embolic protection in saphenous vein graft intervention: results in 49,325 senior patients in the Medicare-linked National Cardiovascular Data Registry Cath PCI Registry. *Circ Cardiovasc Interv* 8:e001403, 2015
- [65] Liu W, Zou Z, Jiang H et al. Clinical effect of preoperative high-dose atorvastatin against no-reflow after PCI. *Exp Ther Med* 13:97–102, 2017
- [66] Wang J, Chen Y, Wang C et al. The impact of pre-primary percutaneous coronary intervention β blocker use on the no-reflow phenomenon in patients with

acute myocardial infarction. *Zhonghua Xin Xue Guan Bing Za Zhi* 42:822–6, 2014

17.3 Koronarperforation

- [1] Cowley MJ, Dorros G, Kelsey SF et al. Acute coronary events associated with percutaneous transluminal coronary angioplasty. *Am J Cardiol* 53:12C–6C, 1984
- [2] Dippel EJ, Kereiakes DJ, Tramuta DA et al. Coronary perforation during percutaneous coronary intervention in the era of abciximab platelet glycoprotein IIb/IIIa blockade: an algorithm for percutaneous management. *Cathet Cardiovasc Interv* 52:279–86, 2001
- [3] Ellis SG, Ajluni S, Arnold AZ et al. Increased coronary perforation in the new device era. Incidence, classification, management, and outcome. *Circulation* 90:2725–30, 1994
- [4] Eggebrecht H, Ritzel A, von Birgelen C et al. Acute and long-term outcome after coronary artery perforation during percutaneous coronary interventions. *Z Kardiol* 93:791–798, 2004
- [5] Gruberg L, Pinnow E, Flood R et al. Incidence, management, and outcome of coronary artery perforation during percutaneous coronary intervention. *Am J Cardiol* 86:680–2, 2000
- [6] Gunning MG, Williams IL, Jewitt DE et al. Coronary artery perforation during percutaneous intervention: incidence and outcome. *Heart* 88:495–8, 2002
- [7] Stankovic G, Orlic D, Corvaja N et al. Incidence, predictors, in-hospital, and late outcome of coronary artery perforations. *Am J Cardiol* 93:213–6, 2004
- [8] Lemmert ME, van Bommel RJ, Diletti R et al. Clinical characteristics and management of coronary artery perforations: a single-center 11-year experience and practical overview. *J Am Heart Assoc* 6:e007049, 2017. <https://www.ahajournals.org/doi/epub/10.1161/JAHA.117.007049>
- [9] Guttman OP, Jones DA, Gulati A et al. Prevalence and outcomes of coronary artery perforation during percutaneous coronary intervention. *EuroIntervention* 13:e595–601, 2017
- [10] Kinnaird T, Anderson R, Ossei-Gerning N et al. Coronary perforation complicating percutaneous coronary intervention in patients with a history of coronary artery bypass surgery: an analysis of 309 perforation cases from the British Cardiovascular Intervention Society Database. *Circ Cardiovasc Interv* 10:e005581, 2017
- [11] Kinnaird T, Shing Kwok C, Kontopantelis E et al. Coronary perforation during PCI between 2006 and 2013: an analysis of 527.121 cases from the BCIS database. *Circ Cardiovasc Interv* 9:e003449, 2016
- [12] Danek BA, Karatasakis A, Tajti P et al. Incidence, treatment, and outcomes of coronary perforation during chronic total occlusion percutaneous coronary intervention. *Am J Cardiol* 120:1285–92, 2017
- [13] Berry JF, von Mering GO, Schmalfuss C et al. Systolic compression of the left anterior descending coronary artery: a case series, review of the literature, and therapeutic options including stenting. *Catheter Cardiovasc Interv* 56:58–63, 2002
- [14] Tomasevic M, Dikic M, Ostojic M. Stenting a myocardial bridge: a wrong decision in STEMI? *Acta Cardiol* 66:89–91, 2011
- [15] Rubio-Alonso B, García-Tejada J, Grandanista C. Coronary perforation to left ventricular cavity following stenting within a myocardial bridge. *Coron Artery Dis* 29:271–3, 2018
- [16] Pourhoseini S, Bakhtiari M, Babaee A et al. Increased risk of coronary perforation during percutaneous intervention of myocardial bridge: what histopathology says. *J Cardiovasc Thorac Res* 9:108–12, 2017. doi: 10.15171/jcvtr.2017.18
- [17] Cortese B, di Palma G, Latini R. Coronary sinus perforation during reducer implantation. *Catheter Cardiovasc Interv* 91:1291–3, 2018
- [18] Mirza AJ, Taha AY, Aldoori JS et al. Coronary artery perforation complicating percutaneous coronary intervention. *Asian Cardiovasc Thorac Ann* 26:101–6, 2018
- [19] Erbel R, O'Neill W, Auth D et al. High-frequency rotablation of occluded coronary artery during heart catheterization. *Cathet Cardiovasc Diagn* 17:56–8, 1989
- [20] Mantis C, Kintis K, Papadakis E et al. Rota-Wire-induced coronary perforation and tamponade during rotational atherectomy. *Kardiol Pol* 76:676, 2018
- [21] Davidson LJ, Ricciardi MJ. Coronary artery perforation complicated by pericardial ab-

- scuss formation: a Clinical Dilemma. *Circ Cardiovasc Interv* 11:e005917, 2018
- [22] Li Y, Xu R, Cai Z et al. Acute purulent pericarditis following staged percutaneous coronary intervention for multivessel disease. *Herz* 38:934–7, 2013
 - [23] Roubelakis A, Rawlins J, Baliulis G. Coronary artery rupture caused by stent infection: a rare complication. *Circulation* 131:1302–3, 2015
 - [24] Fejka M, Dixon SR, Safian RD et al. Diagnosis, management, and clinical outcome of cardiac tamponade complicating percutaneous coronary intervention. *Am J Cardiol* 90:1183–6, 2002
 - [25] Eeckhout E, De Palma R. Coronary perforation: an inconvenient complication. *JACC Cardiovasc Interv* 4:96–7, 2011
 - [26] Colombo A, Itoh A, Di Mario C et al. Successful closure of a coronary vessel rupture with a vein graft stent: case report. *Cathet Cardiovasc Diagn* 38:172–4, 1996
 - [27] Caputo RP, Amin N, Marvasti M et al. Successful treatment of a saphenous vein graft perforation with an autologous vein-covered stent. *Catheter Cardiovasc Interv* 48:382–6, 1999
 - [28] Briguori C, Nishida T, Anzuini A et al. Emergency polytetrafluoroethylene-covered stent implantation to treat coronary ruptures. *Circulation* 102:3028–31, 2000
 - [29] Campbell PG, Hall JA, Harcombe AA et al. The Jomed Covered Stent Graft for coronary artery aneurysms and acute perforation: a successful device which needs careful deployment and may not reduce restenosis. *J Invasive Cardiol* 12:272–6, 2002
 - [30] Dixon SR, Webster MWI, Ormiston JA et al. Gelfoam embolization of a distal coronary artery guidewire perforation. *Cathet Cardiovasc Interv* 49:214–7, 2000
 - [31] George S, Cotton J, Wrigley B. Guidewire-induced coronary perforation successfully treated with subcutaneous fat embolisation: a simple technique available to all. *Catheter Cardiovasc Interv* 86:1186–8, 2015
 - [32] Shemisa K, Karatasakis A, Brilakis ES. Management of guidewire-induced distal coronary perforation using autologous fat particles versus coil embolization. *Catheter Cardiovasc Interv* 89:253–8, 2017
 - [33] Politi L, Iaccarino D, Sangiorgi GM et al. Huge coronary perforation during percutaneous intervention sealed by injection of polyvinyl alcohol microspheres. *J Cardiovasc Med (Hagerstown)* 16(Suppl 2):S130–2, 2015
 - [34] Meincke F, Kuck KH, Bergmann MW. Cardiac tamponade due to coronary perforation during percutaneous interventions successfully treated with microspheres. *Clin Res Cardiol* 103:325–7, 2014
 - [35] Gutiérrez-Chico JL. Zambomballooning: A novel technique to seal a coronary perforation complicating the intervention of a chronic total occlusion. *Cardiol J* 24:573–5, 2017
 - [36] Takagi K, Furui K, Morishima I et al. Usefulness of the jailed-balloon technique in percutaneous intervention for severe coronary perforation involving left main bifurcation. *JACC Cardiovasc Interv* 10:e209–11, 2017
 - [37] Gonzalez-Santos JM, Vallejo JL, Pineda T et al. Emergency surgery after coronary disruption complicating PTCA. Report of four cases. *Thorac Cardiovasc Surg* 33:244–7, 1985
 - [38] Tugtekin SM, Alexiou K, Kappert U et al. Chirurgische Therapie nach traumatischen interventionellen Koronarläsionen. *Z Kardiol* 92:833–6, 2003
 - [39] Karaman K, Karayakali M, Arisoy A et al. A late complication of coronary artery perforation during primary percutaneous coronary intervention: Coronary arteriovenous fistula. *Türk Kardiyol Dern Ars* 45:739–43, 2017
 - [40] Matteucci M, Fina D, Jiritano F et al. Treatment strategies for post-infarction left ventricular free-wall rupture. *Eur Heart J Acute Cardiovasc Care* 8:379–87, 2019
 - [41] Nasrin S, Cader FA, Haq MM et al. Right coronary artery perforation extending to the coronary sinus of Valsalva during percutaneous intervention successfully sealed with polytetrafluoroethylene-covered stent: a case report. *BMC Res Notes* 10:537, 2017
 - [42] Kaneko U, Kashima Y, Kanno D et al. Giant coronary artery pseudoaneurysm formed after septal artery perforation during retrograde percutaneous coronary intervention: a conservative strategy with serial cardiac computed tomographic imaging. *JACC Cardiovasc Interv* 11:e23–4, 2018

- [43] Yamanaka T, Fukatsu T, Ichinohe Y et al. Spontaneous closure of ventricular septal perforation following percutaneous coronary intervention for acute myocardial infarction. *BMJ Case Rep* Oct 13:bcr-221204, 2017
- [44] Kawamoto H, Tanaka K, Ruparelia N et al. Short-term and long-term outcomes after polytetrafluoroethylene-covered stent implantation for the treatment of coronary perforation. *Am J Cardiol* 116:1822–6, 2015

17.4 Periinterventioneller Schlaganfall

- [1] Wong SC, Minutello R, Hong MK. Neurological complications following percutaneous coronary interventions (a report from the 2000–2001 New York State Angioplasty Registry). *Am J Cardiol* 96:1248–50, 2005
- [2] Hamon M, Baron JC, Viader F et al. Periprocedural stroke and cardiac catheterization. *Circulation* 118:678–83, 2008
- [3] Keeley EC, Grines CL. Scraping of aortic debris by coronary guiding catheters: a prospective evaluation of 1,000 cases. *J Am Coll Cardiol* 2:1861–5, 1998
- [4] Eggebrecht H, Haude M, Woertgen U et al. Systematic use of a collagen-based vascular closure device immediately after cardiac catheterization procedures in 1.317 consecutive patients. *Catheter Cardiovasc Interv* 7:486–95, 2002
- [5] Eggebrecht H, Oldenburg O, Dirsch O et al. Potential embolization by atherosclerotic debris dislodged from aortic wall during cardiac catheterization: histological and clinical findings in 7.621 patients. *Catheter Cardiovasc Interv* 49:389–94, 2000
- [5a] Ringleb P, Köhrmann M, Jansen O et al. Akuttherapie des ischämischen Schlaganfalls, S2e-Leitlinie, 2022, in Deutsche Gesellschaft für Neurologie (Hrsg.), Leitlinie für Diagnostik und Therapie in der Neurologie. Online: www.dgn.org/leitlinien, zuletzt bearbeitet 28.11.2022
- [5b] Tužil J, Matějka J, Mamas MA, Doležal T. Short-term risk of periprocedural stroke relative to radial vs. femoral access: systematic review, meta-analysis, study sequential analysis and meta-regression of 2,188,047 real-world cardiac catheterizations. *Expert Rev Cardiovasc Ther* 21:293–304, 2023
- [6] Hamon M, Gomes S, Oppenheim C et al. Cerebral microembolism during cardiac catheterization and risk of acute brain injury: a prospective diffusion-weighted magnetic resonance imaging study. *Stroke* 7:2035–8, 2006
- [7] Kronzon I, Tunick PA. Aortic atherosclerotic disease and stroke. *Circulation* 14:63–75, 2006
- [8] De Marco F, Antonio Fernandez-Diaz J, Lefèvre T et al. Management of cerebrovascular accidents during cardiac catheterization: immediate cerebral angiography versus early neuroimaging strategy. *Catheter Cardiovasc Interv* 70:560–8, 2007
- [9] Al-Mubarak N, Vitek JJ, Mousa I et al. Immediate catheter-based neurovascular rescue for acute stroke complicating coronary procedures. *Am J Cardiol* 90:173–6, 2002
- [10] Janjua N, Brisman JL. Endovascular treatment of acute ischaemic stroke. *Lancet Neurol* 6:1086–93, 2007
- [11] Khatri P, Taylor RA, Palumbo V et al. Treatment of Acute Stroke after Cardiac Catheterization (TASCC) Study Group. The safety and efficacy of thrombolysis for strokes after cardiac catheterization. *J Am Coll Cardiol* 51:906–11, 2008
- [11] Campbell BC, Mitchell PJ, Kleinig TJ et al. Endovascular therapy for ischemic stroke with perfusion-imaging selection. *N Engl J Med*. 372:1009–18, 2015
- [12] Saver JL, Jahan R, Levy EI et al. Solitaire flow restoration device versus the merci retriever in patients with acute ischaemic stroke (SWIFT): a randomised, parallel-group, non-inferiority trial. *Lancet* 380:1241–9, 2012
- [13] Nogueira RG, Lutsep HL, Gupta R et al. Trevo versus Merci retrievers for thrombectomy revascularisation of large vessel occlusions in acute ischaemic stroke (TREVO 2): a randomised trial. *Lancet* 380:1231–40, 2012. Erratum in: *Lancet* 380:1230, 2012
- [14] Berkhemer O, Fransen P, Beumer D et al. A randomized trial of intraarterial treatment for acute ischemic stroke. *N Engl J Med* 372:11–20, 2015

- [15] Jovin TG, Chamorro A, Cobo E et al. Thrombectomy within 8 hours after symptom onset in ischemic stroke. *N Engl J Med* 372, 2296–306, 2015
- [16] Saver JL, Goyal M, Bonafe A et al. Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *N Engl J Med* 372:2285–95, 2015
- [17] Bracard S, Ducrocq X, Mas JL et al. Mechanical thrombectomy after intravenous alteplase versus alteplase alone after stroke (THRACE): a randomised controlled trial. *Lancet Neurol* 15: 1138–1147, 2016
- [18] Nogueira RG, Jadhav AP, Haussen DC et al. Thrombectomy 6 to 24 hours after stroke with a mismatch between deficit and infarct. *N Engl J Med* 378:11–21, 2018
- [19] Saver JL, Goyal M, van der Lugt A et al. Time to treatment with endovascular thrombectomy and outcomes from ischemic stroke: a meta-analysis. *JAMA* 316:1279–88, 2016
- [20] Sheinberg DL, McCarthy DJ, Peterson EC et al. DEFUSE-3 Trial: reinforcing evidence for extended endovascular intervention time window for ischemic stroke. *World Neurosurg* 112:275–6, 2018
- [21] Lavine SD, Cockroft K, Hoh B et al. Training guidelines for endovascular stroke intervention: an international multi-society consensus document. *Neuroradiology* 58:537–41, 2016
- [6] Kieny A, Wettlé C, Lipsker D et al. Earlobe crease and cholesterol emboli. *Ann Dermatol Venereol* 142:619–21, 2015
- [7] Sedano Casas A, Gato Díez A, Mora Escudero I et al. Cholesterol emboli-induced ischemic colitis. *Med Clin (Barc)* 141:e7, 2013
- [8] Shaikh AH, Hanif B, Hasan K et al. Cholesterol emboli syndrome – a rare complication of cardiac catheterization. *J Pak Med Assoc* 60:492–4, 2010
- [9] Frank RD, Velden J. Images in clinical medicine. Cholesterol emboli after coronary bypass surgery. *N Engl J Med* 364:265, 2011
- [10] Pervaiz MH, Durga S, Janoudi A et al. PET/CTA detection of muscle inflammation related to cholesterol crystal emboli without arterial obstruction. *J Nucl Cardiol* 25:433–40, 2018
- [11] Zucchelli P, Zuccalà A. The diagnostic dilemma of hypertensive nephrosclerosis: the nephrologist's view. *Am J Kidney Dis* 21(5 Suppl 2):87–91, 1993
- [12] Scolari F, Ravani P, Pola A et al. Predictors of renal and patient outcomes in attheroembolic renal disease: a prospective study. *J Am Soc Nephrol* 4:1584–90, 2003
- [13] Bashore TM, Gehrig T. Cholesterol emboli after invasive procedures. *J Am Coll Cardiol* 42:217–8, 2003
- [14] Elinav E, Chajek-Shaul T, Stern M. Improvement in cholesterol emboli syndrome after iloprost therapy. *BMJ* 24:268–9, 2002
- [15] Koga J, Ohno M, Okamoto K et al. Cholesterol embolization treated with corticosteroids – two case reports. *Angiology* 6:497–501, 2005
- [16] Simon R, Berwert L, Burnier M et al. Severe renal insufficiency secondary to renal cholesterol emboli: therapeutical options revisited. *Rev Med Suisse* 6:432–4, 436–7, 2010

17.5 Cholesterinemboliesyndrom

- [1] Lusco MA, Najafian B, Alpers CE et al. AJKD atlas of renal pathology: cholesterol emboli. *Am J Kidney Dis* 6:e23–4, 2016
- [2] Rupprecht H. Postinterventionelles Cholesterinemboliesyndrom. *Dtsch Med Wochenschr* 32:1833–36, 2007
- [3] Fine MJ, Kapoor W, Falanga V. Cholesterol crystal embolization: a review of 221 cases in the English literature. *Angiology* 8:769–84, 1987
- [4] Ogbechie OA, Morley KW, Sugai D et al. Limited cutaneous pseudovasculitis: a mild variant of cholesterol emboli syndrome. *Circulation* 131:514–5, 2015
- [5] Rousseau A, de Monchy I, Barreau E et al. Retinal emboli in cholesterol crystal embolism. *Case Rep Ophthalmol Med* 2013:421352, 2013

17.6 Koronarembolie

- [1] Shibata T, Kawakami S, Noguchi T et al. Prevalence, clinical features, and prognosis of acute myocardial infarction attributable to coronary artery embolism. *Circulation* 132:241–50, 2015
- [2] Raphael CE, Heit JA, Reeder GS et al. Coronary embolus: an underappreciated

- cause of acute coronary syndromes. *JACC Cardiovasc Interv* 11:172–80, 2018
- [3] Jungbluth A, Erbel R, Darius H et al. Paradoxical coronary embolism: case report and review of the literature. *Am Heart J* 116:879–85, 1988
 - [4] Marchand F. Zur Kenntnis der Embolie und Thrombose der Gehirnarterien, zugleich ein Beitrag zur Casuistik der primären Herztumoren und der gekreuzten Embolie. *Berl Klin Wochenschr* 31:36, 1894
 - [5] Wąsek WC, Samul W, Ryzek R et al. Unique case of ST-segment-elevation myocardial infarction related to paradoxical embolization and simultaneous pulmonary embolization: clinical considerations on indications for patent foramen ovale closure in no-guidelines land. *Circulation* 131:1214–23, 2015
 - [6] Mügge A, Daniel WG, Wenzlaff P et al. Patent foramen ovale or left atrial thrombi in unexplained arterial embolism. *Lancet* 1:282–3, 1989
 - [7] Hausmann D, Mügge A, Becht I et al. Diagnosis of patent foramen ovale by transesophageal echocardiography and association with cerebral and peripheral embolic events. *Am J Cardiol* 70:668–72, 1992
 - [8] Hausmann D, Mügge A, Daniel WG. Identification of patent foramen ovale permitting paradoxical embolism. *J Am Coll Cardiol* 26:1030–8, 1995
 - [9] Nellessen U, Daniel WG, Matheis G et al. Impending paradoxical embolism from atrial thrombus: correct diagnosis by transesophageal echocardiography and prevention by surgery. *J Am Coll Cardiol* 5:1002–4, 1985
 - [10] Fauveau E, Cohen A, Bonnet N et al. Surgical or medical treatment for thrombus straddling the patent foramen ovale: impending paradoxical embolism? Report of four clinical cases and literature review. *Arch Cardiovasc Dis* 101:637–44, 2008
 - [11] Wahl A, Jüni P, Mono ML et al. Long-term propensity score-matched comparison of percutaneous closure of patent foramen ovale with medical treatment after paradoxical embolism. *Circulation* 125:803–12, 2012
 - [12] Uchida S, Yamamoto M, Masaoka Y et al. A case of acute pulmonary embolism and acute myocardial infarction with suspected paradoxical embolism after laparoscopic surgery. *Heart Vessels* 14:197–200, 1999
 - [13] Haghi D, Sueselbeck T, Papavassiliou T et al. Paradoxical coronary embolism causing non-ST segment elevation myocardial infarction in a case of pulmonary embolism. *Z Kardiol* 93:824–8, 2004
 - [14] Rovner A, Valika AA, Kovacs A et al. Possible paradoxical embolism as a rare cause for an acute myocardial infarction. *Echocardiography* 23:407–9, 2006
 - [15] Konstantinides S, Geibel A, Kasper W et al. Patent foramen ovale is an important predictor of adverse outcome in patients with major pulmonary embolism. *Circulation* 97:1946–51, 1998
 - [16] Kälsch H, Wieneke H, Erbel R. Acute myocardial infarction in a patient with chronic myelocytic leukemia during chemotherapy with hydroxyurea. *Herz* 35:420–2, 2010
 - [17] Gutersohn A, Zimmermann U, Bartel T et al. A rare case of acute „infective“ myocardial infarction triggered by acute parvovirus B19 myocarditis. *Nat Clin Pract Cardiovasc Med* 2:167–71, 2005
 - [18] Bommer M, Wölfle-Guter M, Bohl S et al. The differential diagnosis and treatment of thrombotic microangiopathies. *Dtsch Arztebl Int* 115:327–34, 2018
 - [19] Patschan D, Witzke O, Dührsen U et al. Acute myocardial infarction in thrombotic microangiopathies – clinical characteristics, risk factors and outcome. *Nephrol Dial Transplant* 21:1549–54, 2006
 - [20] Brunson JG. Coronary embolism in bacterial endocarditis. *Am J Pathol* 29:689–701, 1953
 - [21] Erbel R, Liu F, Ge J et al. Identification of high-risk subgroups in infective endocarditis and the role of echocardiography. *Eur Heart J* 16:588–602, 1995
 - [22] Breuckmann F, Naber CK, Boese D et al. Mitral valve endocarditis leading to acute myocardial and cerebellar infarction in a young adult. *Clin Res Cardiol* 95:657–62, 2006
 - [23] Dauvergne C, Araya M, Valenzuela J et al. Acute myocardial infarction after left-heart catheterization in a patient with severe calcified bicuspid aortic stenosis. *JACC Cardiovasc Interv* 7:e5–6, 2014
 - [24] Murai T, Yonetsu T, Isobe M et al. Coronary embolization caused by pleomorphic

- lung carcinoma. *Intern Med* 55:3607–9, 2016
- [24] Cheng TO. Coronary embolism. *Int J Cardiol* 136:1–3, 2009
- [25] O'Neill W, Timmis GC, Bourdillon PD et al. A prospective randomized clinical trial of intracoronary streptokinase versus coronary angioplasty for acute myocardial infarction. *N Engl J Med* 314:812–8, 1986
- [26] Nunn CM, O'Neill WW, Rothbaum D et al. Long-term outcome after primary angioplasty: report from the primary angioplasty in myocardial infarction (PAMI-I) trial. *J Am Coll Cardiol* 33:640–6, 1999
- [27] Mehta RH, Harjai KJ, Cox DA et al. Comparison of coronary stenting versus conventional balloon angioplasty on five-year mortality in patients with acute myocardial infarction undergoing primary percutaneous coronary intervention. *Am J Cardiol* 96:901–6, 2005
- [28] Kunadian B, Dunning J, Vijayalakshmi K et al. Meta-analysis of randomized trials comparing anti-embolic devices with standard PCI for improving myocardial reperfusion in patients with acute myocardial infarction. *Catheter Cardiovasc Interv* 69:488–96, 2007
- [29] Kuntz RE, Rogers C, Baim DS. Percutaneous coronary intervention-induced emboli during primary PCI for STEMI: too little, too much, or too late? *Am Heart J* 150:4–6, 2005
- [30] Brown ED, Blankenship JC. A mechanism for stroke complicating thrombus aspiration. *Catheter Cardiovasc Interv* 89:93–6, 2017
- [31] Eggebrecht H, Baumgart D, Naber C et al. Extraction of large intracoronary thrombus in acute myocardial infarction by percutaneous Fogarty maneuver: intentional abuse of a novel interventional device. *Catheter Cardiovasc Interv* 55:228–32, 2002
- [32] Satogami K, Ino Y, Kubo T et al. Impact of plaque rupture detected by optical coherence tomography on transmural extent of infarction after successful stenting in ST-segment elevation acute myocardial infarction. *JACC Cardiovasc Interv* 10:1025–33, 2017
- [33] Eick B, Haude M, Altmann C et al. Treatment of a large intracoronary thrombus with urokinase and a chimeric monoclonal platelet aggregation inhibitor. *Dtsch Med Wochenschr* 122:709–15, 1997
- [34] Kahles T, Garcia-Esperon C, Zeller S et al. Mechanical thrombectomy using the new ERIC retrieval device is feasible, efficient, and safe in acute ischemic stroke: A Swiss Stroke Center Experience. *Am J Neuroradiol* 37:114–9, 2016
- [35] Manchurov V, Anisimov K, Khokhlova V et al. The first reported stentriever-based thrombectomy in acute ST-elevation myocardial infarction due to paradoxical coronary embolism. *Eurointerv* 13:e602–3, 2017
- [36] Popovic B, Agrinier N, Bouchahda N et al. Coronary embolism among ST-segment-elevation myocardial infarction patients: mechanisms and management. *Circ Cardiovasc Interv* 11:e005587, 2018

17.7 Bergung von Fremdkörpern

- [1] Alkhouli M, Sievert H et al. Device embolization in structural heart interventions: incidence, outcomes and retrieval techniques. *JACC Cardiovasc Interv* 12:113–26, 2019
- [2] Veldhuijzen FL, Michels HR, el Gamal MI et al. Retrieval of undeployed stents from the right coronary artery: report of two cases. *Cathet Cardiovasc Diagn* 30:245–8, 1993